

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **c-Myc Antibody, Clone: [MYC909], Mouse, Monoclonal NSJ-V2746IHC-7ML**

Article Name	c-Myc Antibody, Clone: [MYC909], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2746IHC-7ML
Supplier Catalog Number	V2746IHC-7ML
Alternative Catalog Number	NSJ-V2746IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the c-Myc antibody.
Product Description	It recognizes a transcription factor of 64-67kDa, identified as c-myc. This mAb shows no cross-reaction with v-myc. c-Myc is a transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5-CAC[GA]T...
Clonality	Monoclonal
Clone Designation	[MYC909]
UniProt	<a href="#">P01106</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	The stated application concentrations are suggested starting amounts. Optimal dilution of the c-Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.