

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal NSJ-V2747-100UG

| | |
|----------------------------|---|
| Article Name | c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V2747-100UG |
| Supplier Catalog Number | V2747-100UG |
| Alternative Catalog Number | NSJ-V2747-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC-P |
| Species Reactivity | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the c-Myc antibody cocktail. |
| Product Description | It recognizes a transcription factor of 64-67kDa, identified as c-myc. This mAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a variety of tumo... |
| Clonality | Monoclonal |
| Clone Designation | [MYC275 + MYC909] |
| UniProt | P01106 |
| Purity | Protein G affinity chromatography |

| | |
|--------------------|---|
| Form | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | Optimal dilution of the c-Myc antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min |