

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal NSJ-V2747-20UG**

Article Name	c-Myc Antibody Cocktail, Clone: [MYC275 + MYC909], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2747-20UG
Supplier Catalog Number	V2747-20UG
Alternative Catalog Number	NSJ-V2747-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the c-Myc antibody cocktail.
Product Description	It recognizes a transcription factor of 64-67kDa, identified as c-myc. This mAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a variety of tumo...
Clonality	Monoclonal
Clone Designation	[MYC275 + MYC909]
UniProt	<a href="#">P01106</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the c-Myc antibody cocktail should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min