

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### NCAM Antibody, Clone: [NCAM1/784], Mouse, Monoclonal NSJ-V2753-100UG

Article Name	NCAM Antibody, Clone: [NCAM1/784], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2753-100UG
Supplier Catalog Number	V2753-100UG
Alternative Catalog Number	NSJ-V2753-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the NCAM antibody.
Product Description	This mAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic...
Clonality	Monoclonal
Clone Designation	[NCAM1/784]
UniProt	<a href="#">P13591</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Application Notes	Optimal dilution of the NCAM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.