

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

NCL Antibody / Nucleolin, Clone: [364-5], Mouse, Monoclonal NSJ-V2757IHC-7ML

Article Name	NCL Antibody / Nucleolin, Clone: [364-5], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2757IHC-7ML
Supplier Catalog Number	V2757IHC-7ML
Alternative Catalog Number	NSJ-V2757IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	SUDHL1 cell nuclear lysate was used as the immunogen for the NCL antibody.
Product Description	Recognizes a protein of ~76kDa, which is identified as Nucleolin (NCL). It is the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleola...
Clonality	Monoclonal
Clone Designation	[364-5]
UniProt	P19338
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the NCL antibody should be determined by the researcher.1. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.