

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### NuMA Antibody, Clone: [SPM300], Mouse, Monoclonal NSJ-V2768-20UG

|                            |   |
|----------------------------|---|
| Article Name               | NuMA Antibody, Clone: [SPM300], Mouse, Monoclonal   |
| Biozol Catalog Number      | NSJ-V2768-20UG  |
| Supplier Catalog Number    | V2768-20UG  |
| Alternative Catalog Number | NSJ-V2768-20UG  |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC-P   |
| Species Reactivity         | Human   |
| Immunogen                  | Colon carcinoma 174T cells were used as the immunogen for the NuMA antibody.  |
| Product Description        | Recognizes a phosphorylated protein of 228kDa, identified as nuclear mitotic apparatus protein (NuMA). NuMA is intra-nuclear protein and present in nucleus during interphase. At the onset of mitosis, it redistributes from the nucleus to two centrosom... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [SPM300]  |
| UniProt                    | <a href="#">Q14980</a>  |
| Purity                     | PEG precipitation   |

|                    |   |
|--------------------|---|
| Form               | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide  |
| Antibody Type      | Primary Antibody  |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT  |
| Application Notes  | Optimal dilution of the NuMA antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min. |