

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD31 Antibody Cocktail, Clone: [C31.3 + C31.7 + C31.10], Mouse, Monoclonal NSJ-V2784-20UG

Article Name	CD31 Antibody Cocktail, Clone: [C31.3 + C31.7 + C31.10], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2784-20UG
Supplier Catalog Number	V2784-20UG
Alternative Catalog Number	NSJ-V2784-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P, WB
Species Reactivity	Human
Immunogen	Human recombinant protein was used as the immunogen for the CD31 antibody cocktail.
Product Description	CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime...
Clonality	Monoclonal
Clone Designation	[C31.3 + C31.7 + C31.10]
UniProt	P16284
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the CD31 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.