

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Cytochrome C Antibody, Clone: [SPM389], Mouse, Monoclonal NSJ-V2787IHC-7ML

Article Name	Cytochrome C Antibody, Clone: [SPM389], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2787IHC-7ML
Supplier Catalog Number	V2787IHC-7ML
Alternative Catalog Number	NSJ-V2787IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids 66-104 of pigeon Cytochrome C were used as the immunogen for the Cytochrome C antibody. It recognizes an epitope within amino acids 93-104 of pigeon Cytochrome C.
Product Description	Cytochrome C is a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is normally localized to the mitochondrial inter-membrane space....
Clonality	Monoclonal
Clone Designation	[SPM389]
UniProt	<a href="#">P99999</a>
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the Cytochrome C antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.