

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant p40 Antibody / deltaNp63, Clone: [rTP40/3690], Biotin, Mouse, Monoclonal NSJ-V8627BTN

| | |
|----------------------------|---|
| Article Name | Recombinant p40 Antibody / deltaNp63, Clone: [rTP40/3690], Biotin, Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V8627BTN |
| Supplier Catalog Number | V8627BTN |
| Alternative Catalog Number | NSJ-V8627BTN |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC-P |
| Species Reactivity | Human |
| Immunogen | A synthetic peptide from the N-terminal of human p40 protein was used as the immunogen for the recombinant p40 antibody. |
| Conjugation | Biotin |
| Product Description | p63 consists of two major isoforms-TAp63 and delta-Np63. These isoforms differ in the structure of the N-terminal domains. The TAp63 isoform (identified by anti-p63 antibody) contains a transactivation-competent TA domain with homology to p53, which ... |
| Clonality | Monoclonal |
| Clone Designation | [rTP40/3690] |
| UniProt | Q9H3D4 |

| | |
|--------------------|---|
| Purity | Protein G affinity chromatography |
| Form | 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 2-4ug/million cells, Immunofluorescence: 2-4ug/ml, Immunohistochemistry (FFPE): 2-4ug/ml for 30 minutes at RT |
| Application Notes | Optimal dilution of the recombinant p40 antibody should be determined by the researcher. |