

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal NSJ-V9102-20UG**

Article Name	Bcl10 Antibody, Clone: [SPM520], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V9102-20UG
Supplier Catalog Number	V9102-20UG
Alternative Catalog Number	NSJ-V9102-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids 122-168 were used as the immunogen for this Bcl10 antibody.
Product Description	Bcl10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1,14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of Bcl10...
Clonality	Monoclonal
Clone Designation	[SPM520]
UniProt	<a href="#">O95999</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)
Application Notes	<p>The optimal dilution of the Bcl10 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>