

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Chromogranin A Antibody, Clone: [CHGA/798], Unconjugated, Mouse, Monoclonal BYT-ORB2638303

| | |
|----------------------------|---|
| Article Name | Chromogranin A Antibody, Clone: [CHGA/798], Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | BYT-ORB2638303 |
| Supplier Catalog Number | orb2638303 |
| Alternative Catalog Number | BYT-ORB2638303-100 |
| Manufacturer | Biorbyt |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant human protein was used as the immunogen for the Chromogranin A antibody. |
| Conjugation | Unconjugated |
| Product Description | Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para... |
| Clonality | Monoclonal |
| Clone Designation | [CHGA/798] |
| UniProt | P10645 |

| | |
|--------------------|---|
| Buffer | 1 mg/ml in 1X PBS, rAlbumin free, sodium azide free |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | <p>Application Notes: Optimal dilution of the Chromogranin A antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p> |