

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

EpCAM Antibody / Cytoplasmic domain, Clone: [EGP40/1110], Unconjugated, Mouse, Monoclonal BYT-ORB2638811

Article Name	EpCAM Antibody / Cytoplasmic domain, Clone: [EGP40/1110], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB2638811
Supplier Catalog Number	orb2638811
Alternative Catalog Number	BYT-ORB2638811-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Canine, Feline, Human, Mouse, Rat
Immunogen	A portion of amino acids 289-314 (cytoplasmic domain) was used as the immunogen for the EpCAM antibody.
Conjugation	Unconjugated
Product Description	EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast major...
Clonality	Monoclonal
Clone Designation	[EGP40/1110]
UniProt	P16422

Buffer	1 mg/ml in 1X PBS, rAlbumin free, sodium azide free
Application Dilute	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Application Notes	Application Notes: Optimal dilution of the EpCAM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min