

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal BYT-ORB605908

Article Name	MUC1 Antibody / Mucin-1 / EMA, Clone: [MUC1/520], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB605908
Supplier Catalog Number	orb605908
Alternative Catalog Number	BYT-ORB605908-100,BYT-ORB605908-20
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	IHC-P, WB
Species Reactivity	Human
Immunogen	Human milk-fat globule membranes (HMFGM) was used as the immunogen for the MUC1 antibody.
Conjugation	Unconjugated
Product Description	In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of MUC1/Mucin-1/Epithelial Marker Antigen/EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhes...
Clonality	Monoclonal
Clone Designation	[MUC1/520]
UniProt	P15941

Buffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Western blot: 1-2ug/ml
Application Notes	Application Notes: Optimal dilution of the MUC1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min