

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

H1F0 Antibody, Clone: [1415-1], Unconjugated, Mouse, Monoclonal BYT-ORB1252534

Article Name	H1F0 Antibody, Clone: [1415-1], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB1252534
Supplier Catalog Number	orb1252534
Alternative Catalog Number	BYT-ORB1252534-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC-P, WB
Species Reactivity	Human, Mouse, Rat
Immunogen	Nuclei of human leukemia biopsy cells were used as the immunogen for the Histone antibody.
Conjugation	Unconjugated
Product Description	H1F0 Antibody...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[1415-1]
UniProt	P07305
Buffer	PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Form	Liquid

Target	H1F0
Application Notes	<p>Application Notes: Western blot: 0.5-1 µg/mL Flow Cytometry: 0.5-1 µg/million cells in 0.1mL Immunofluorescence: 0.5-1 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 min at RT (1) Prediluted format : incubate for 30 min at RT (2) Optimal dilution of the Histone antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p>