

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PTPRC Antibody, Clone: [158-4D3], Unconjugated, Mouse, Monoclonal BYT-ORB1252710

| | |
|----------------------------|--|
| Article Name | PTPRC Antibody, Clone: [158-4D3], Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | BYT-ORB1252710 |
| Supplier Catalog Number | orb1252710 |
| Alternative Catalog Number | BYT-ORB1252710-100 |
| Manufacturer | Biorbyt |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Stimulated human leukocytes were used as the immunogen for this CD45RA antibody. |
| Conjugation | Unconjugated |
| Product Description | PTPRC Antibody... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [158-4D3] |
| UniProt | P08575 |
| Buffer | PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide |

| | |
|-------------------|--|
| Form | Liquid |
| Target | PTPRC |
| Application Notes | <p>Application Notes: Flow Cytometry: 0.5-1 ug/million cells IF: 1-2 µg/mL IHC (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2) The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD45RA antibody to be titrated up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p> |