

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD43 Antibody, Clone: [SPN/839], Unconjugated, Mouse, Monoclonal BYT-ORB749794

Article Name	CD43 Antibody, Clone: [SPN/839], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB749794
Supplier Catalog Number	orb749794
Alternative Catalog Number	BYT-ORB749794-20,BYT-ORB749794-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the CD43 antibody.
Conjugation	Unconjugated
Product Description	It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-...
Clonality	Monoclonal
Clone Designation	[SPN/839]
UniProt	<a href="#">P16150</a>

Buffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Application Dilute	Flow cytometry: 1-2ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Application Notes: Optimal dilution of the CD43 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min