

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Phospho-PLK1 (Thr210) (Clone: C2) rabbit mAb APC conjugate, Clone: [PLK1T210-C2], Rabbit, Monoclonal ABI-12-4280-10

| | |
|----------------------------|---|
| Article Name | Phospho-PLK1 (Thr210) (Clone: C2) rabbit mAb APC conjugate, Clone: [PLK1T210-C2], Rabbit, Monoclonal |
| Biozol Catalog Number | ABI-12-4280-10 |
| Supplier Catalog Number | 12-4280-10 |
| Alternative Catalog Number | ABI-12-4280-10-10TESTS |
| Manufacturer | Abeomics |
| Host | Rabbit |
| Category | Antikörper |
| Application | FACS |
| Species Reactivity | Human |
| Immunogen | A synthetic phospho-peptide corresponding to residues surrounding Thr210 of human phospho PLK1 |
| Conjugation | APC |
| Product Description | PLK1 is activated by phosphorylation of Thr210 in its activation loop. When dephosphorylated, PLK1 auto-inhibition occurs when the Polo-box domain (PDB) binds (and thus inhibits) the kinase domain. PLK1 phosphorylation is directly linked to the cell ... |
| Clonality | Monoclonal |
| Clone Designation | [PLK1T210-C2] |
| Form | Conjugated |

Application Dilute

For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells or 5 μL per 100 μL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.