

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Monoclonal Antibody to IPO-38 (Proliferation Marker)(Clone : SPM260), Clone: [SPM260], Mouse ABI-36-11054**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Monoclonal Antibody to IPO-38 (Proliferation Marker)(Clone : SPM260), Clone: [SPM260], Mouse                                                                                                                                                                  |
| Biozol Catalog Number      | ABI-36-11054                                                                                                                                                                                                                                                  |
| Supplier Catalog Number    | 36-11054                                                                                                                                                                                                                                                      |
| Alternative Catalog Number | ABI-36-11054-100UG                                                                                                                                                                                                                                            |
| Manufacturer               | Abeomics                                                                                                                                                                                                                                                      |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | FACS, IF, IHC                                                                                                                                                                                                                                                 |
| Species Reactivity         | Human, Mouse, Rat                                                                                                                                                                                                                                             |
| Immunogen                  | Spleen cells of a patient with hairy cell leukemia                                                                                                                                                                                                            |
| Product Description        | Recognizes a protein of 14-16kDa, which is a novel nuclear antigen of proliferating cells. IPO-38 antigen is present in the nuclei of proliferating cells such as Hodgkin s disease and non-Hodgkin s lymphomas, different forms of leukemias, breast and ... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Clone Designation          | [SPM260]                                                                                                                                                                                                                                                      |
| Purity                     | Affinity Chromatography                                                                                                                                                                                                                                       |
| Form                       | Purified                                                                                                                                                                                                                                                      |

|                    |                                                                                                                                                                                                                                                                 |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target             | IPO-38 (Proliferation Marker)                                                                                                                                                                                                                                   |
| Application Dilute | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min a |