

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Monoclonal Antibody to Myeloid Cell Marker (Macrophage / Granulocyte Marker)(Clone : BM-2), Clone: [BM-2], Mouse ABI-36-11072

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Monoclonal Antibody to Myeloid Cell Marker (Macrophage / Granulocyte Marker)(Clone : BM-2), Clone: [BM-2], Mouse                                                                                                                                              |
| Biozol Catalog Number      | ABI-36-11072                                                                                                                                                                                                                                                  |
| Supplier Catalog Number    | 36-11072                                                                                                                                                                                                                                                      |
| Alternative Catalog Number | ABI-36-11072-100UG                                                                                                                                                                                                                                            |
| Manufacturer               | Abeomics                                                                                                                                                                                                                                                      |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | FACS, IF, IHC                                                                                                                                                                                                                                                 |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes                                                                                                                                                                                    |
| Product Description        | Recognizes 183kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Clone Designation          | [BM-2]                                                                                                                                                                                                                                                        |
| Purity                     | Affinity Chromatography                                                                                                                                                                                                                                       |
| Form                       | Purified                                                                                                                                                                                                                                                      |

|                    |                                                                                                                                                                                                                                                                |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target             | Myeloid Cell Marker (Macrophage / Granulocyte Marker)                                                                                                                                                                                                          |
| Application Dilute | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min |