

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rat anti Mouse alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal NMB-RAAM/A2M

| | |
|----------------------------|---|
| Article Name | Rat anti Mouse alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RAAM/A2M |
| Supplier Catalog Number | RaAM/A2M |
| Alternative Catalog Number | NMB-RAAM/A2M |
| Manufacturer | NordicMubio |
| Host | Rat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to alpha-2 macroglobulin as tested in immunoelectrophoresis and radial immunodiffusion. A characteristic single precipitin line is obtained with normal serum which shows a reaction of identity with the pr... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rat serum. No foreign proteins added. Reconstitute the lyophilized ant |
| Source | Alpha-2 macroglobulin ids the major protein in the alpha-2 region of mouse serum after separation by electrophoresis. It is a large polymer and a strong antigen. It forms about one-third in the total of the alpha-2 globulins. The protein is isolated by am |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rat serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. |