

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Swine anti Human IgG (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHU/IGG(FC)

| | |
|----------------------------|---|
| Article Name | Swine anti Human IgG (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal |
| Biozol Catalog Number | NMB-SWAHU/IGG(FC) |
| Supplier Catalog Number | SwAHu/IgG(Fc) |
| Alternative Catalog Number | NMB-SWAHU/IGG(FC) |
| Manufacturer | NordicMubio |
| Host | Porcine |
| Category | Antikörper |
| Species Reactivity | Human |
| Conjugation | Unconjugated |
| Format | IgG |
| Target Specificity | IgG (Fc) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. In immunoelectrophoresis and double radial immunodiffusion, using various antiserum concentrations against normal human plasma and serum, a characteristic precipitin I... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized |
| Source | Purified normal IgG isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l in a 3 mm diameter center well and 2 μ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. |