

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (F(ab)2)-Biotin, MinX Hu DNA-SEC-183399

Article Name	Goat IgG anti-Rabbit IgG (F(ab)2)-Biotin, MinX Hu
Biozol Catalog Number	DNA-SEC-183399
Supplier Catalog Number	SEC-183399
Alternative Catalog Number	DNA-SEC-183399
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rabbit
Immunogen	Rabbit IgG F(ab)2 fragment
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	Human
Product Description	Anti-Rabbit IgG F(ab)2 Antibody generated in goat recognizes the dimeric Fab portion of the rabbit IgG molecule. Rabbit IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controll...
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab') ₂ and Rabbit Serum. No reaction was observed against Rabbit IgG F(c) or Human Serum Proteins.
Form	Lyophilized
Formula	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN ₃
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:12,000 to 1:50,000 of the reconstitution concentration is suggested for this product.