

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (H+L)-Alk. Phos., MinX none DNA-SEC-183852

Article Name	Goat F(ab)2 anti-Rat IgG (H+L)-Alk. Phos., MinX none
Biozol Catalog Number	DNA-SEC-183852
Supplier Catalog Number	SEC-183852
Alternative Catalog Number	DNA-SEC-183852
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rat
Immunogen	Rat IgG whole molecule
Conjugation	Alk. Phos.
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Rat IgG Antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. F(ab)2 Anti-Rat IgG Antibody is ideal for investigators in Cancer, Immunology, and Micr...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:4,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	F(ab') ₂ Anti-Rat IgG Antibody has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for F(ab') ₂ Anti-Rat IgG Antibody.